

Multiple polylogarithm values at roots of unity

Jianqiang Zhao

Department of Mathematics, Eckerd College, St. Petersburg, FL 33711, USA

Abstract. For any positive integer N let μ_N be the group of the N th roots of unity. In this note we shall study the $\mathbb{Q}$ -linear relations among values of multiple polylogarithms evaluated at μ_N . We show that the standard relations considered by Racinet do not provide all the possible relations in the following cases: (i) level $N = 4$, weight $w = 3$ or 4 , and (ii) $w = 2$, $7 < N < 50$, and N is a power of 2 or 3, or N has at least two prime factors. We further find some (presumably all) of the missing relations in (i) by using the octahedral symmetry of $\mathbb{P}^1 - (\{0, \infty\} \cup \mu_4)$. We also prove some other results when $N = p$ or $N = p^2$ (p prime ≥ 5) by using the motivic fundamental group of $\mathbb{P}^1 - (\{0, \infty\} \cup \mu_N)$.

1 Introduction

Double shuffle relations have played significant roles in the study of multiple zeta values in recent years. These relations can be easily generalized to the special values of multiple polylogarithms at roots of unity (MPV for short):

$$Li_{s_1, \dots, s_n}(\zeta_1, \dots, \zeta_n) := \sum_{k_1 > \dots > k_n > 0} \frac{\zeta_1^{k_1} \dots \zeta_n^{k_n}}{k_1^{s_1} \dots k_n^{s_n}}, \quad (s_1, \zeta_1) \neq (1, 1), \quad (1)$$

where ζ_j are all N th roots of unity for $j = 1, \dots, n$. We call N the *level* and $w := s_1 + \dots + s_n$ the *weight*. One of our major interests is to find the dimension $d(w, N)$ of the $\mathbb{Q}$ -vector space $\mathcal{MPV}(w, N)$ spanned by these values.

In [7], Racinet listed the following relations: double shuffle relations and their regularized versions (by regularizing both the divergent integrals and divergent series representing MPVs), (regularized) distributions, and weight one relations. We will call these *standard relations*. By intensive MAPLE computation it is shown [8] that in many cases these relations cannot produce all the possible $\mathbb{Q}$ -linear relations. For example, standard relations imply only $d(3, 4) \leq 9$ which is one more than the bound given by [4, 5.25]. Concretely, from standard relations MAPLE confirms the following

$$\begin{aligned} \text{Fact: Every MPV of weight 3 and level 4 can be written explicitly} \\ \text{as a linear combination of the nine MPVs appearing in (2),} \end{aligned} \quad (*)$$

and no further $\mathbb{Q}$ -linear relations between these values can be deduced from the standard relations. But by GiNac [6] and EZface [1] the following is found numerically (see [8, Remark 10.1])

$$\begin{aligned} 5Li_{1,2}(-1, -i) = 46Li_{1,1,1}(i, 1, 1) - 7Li_{1,1,1}(-1, -1, i) - 13Li_{1,1,1}(i, i, i) + 13Li_{1,2}(-i, i) \\ - Li_{1,1,1}(-i, -1, 1) + 25Li_{1,1,1}(-i, 1, 1) - 8Li_{1,1,1}(i, i, -1) + 18Li_{2,1}(-i, 1). \end{aligned} \quad (2)$$

In this note, we shall prove (2) from the explicit relations in (*) by using the octahedral symmetry of $\mathbb{P}^1 - (\{0, \infty\} \cup \mu_4)$ where μ_N denotes the set of N th roots of unity. We can treat the weight 4 and level 4 case in a similar fashion.

The result above in level four case shows that the standard relations are not always sufficient to determine $d(w, N)$. On the other hand, Deligne and Goncharov [4] find some closed formulae for upper bounds of $d(w, N)$ for all w and N by studying the motivic fundamental group of $\mathbb{P}^1 - (\{0, \infty\} \cup \mu_N)$ in the framework of mixed Tate motives over the ring of S -integers of a number field.

Proposition 1. ([4, 5.25]) *Let N be a positive integer. Let φ be the Euler's totient function and $\nu(N)$ be the number of prime factors of N . Then $d(w, N) \leq D(w, N)$ where $D(w, N)$ are defined by the formal power series*

$$1 + \sum_{w=1}^{\infty} D(w, N)t^w = \begin{cases} (1 - t^2 - t^3)^{-1}, & \text{if } N = 1; \\ (1 - t - t^2)^{-1}, & \text{if } N = 2; \\ (1 - (\frac{\varphi(N)}{2} + \nu(N))t + (\nu(N) - 1)t^2)^{-1}, & \text{if } N \geq 3. \end{cases}$$

As an example one can find the bound $d(w, 4) \leq 2^w$ for all weight w and by a variant of a conjecture of Grothendieck it can be deduced that for $N = 4$ equality should always hold. Further, Goncharov [4] has defined some pro-nilpotent Lie algebra D graded by weight and depth using standard relations and then related D to the motivic fundamental Lie algebra in lower depth cases. Using this we shall improve the bounds in the above proposition for all levels $N = p$ and $N = p^2$ ($p \geq 5$ a prime) in the last section. From this improved bound we find that when $7 < N < 50$ standard relations are incomplete if N is a power of 2 or 3, or N has at least two prime factors, though we don't know how to find missing relations explicitly.

2 The motivic setup

Let us review some constructions from [4]. Fix a positive integer N . Let k be the field over $\mathbb{Q}$ generated by a primitive N th root of unity and $\mathcal{O}$ its ring of integers. Let Γ be the $\mathbb{Q}$ -vector subspace of $\mathcal{O}[1/N]^* \otimes \mathbb{Q}$ generated by $1 - \zeta$ for ζ an N th root of unity other than 1. We refer to [4] for the definition of the tannakian category of mixed Tate motives $\text{MT}(k)_{\Gamma}$, the definition of the fibre functor $\omega(M) := \bigoplus_n \text{Hom}(\mathbb{Q}(n), \text{Gr}_{-2n}^W(M))$, and of the corresponding motivic fundamental group G . Let $\mathcal{L}$ be the Lie algebra generated by generators $e_x, x \in \{0, \infty\} \cup \mu_N$, with the only relation $\sum e_x = 0$. Set the degree of e_x to be 1. Then $\mathcal{L}$ is freely generated by the $e_x, x \neq \infty$. Let Π be the pro-nilpotent group $\lim_n \exp(\mathcal{L}/\deg^{>n})$. The completed enveloping algebra of $\mathcal{L}$ is $\mathbb{Q}\langle\langle e_{\zeta} \rangle\rangle_{\zeta \in \mu_N \cup \{0\}}$ with the coproduct $\Delta : e_{\zeta} \mapsto e_{\zeta} \otimes 1 + 1 \otimes e_{\zeta}$, and Π is the subscheme of the group-like elements under Δ .

Let H_{ω} be the group of μ_N -equivariant automorphisms of Π defined in [4, §5.8]. Set $G_{\omega} := \omega(G)$, $U_{\omega} := \ker(G_{\omega} \rightarrow \mathbb{G}_m)$ and $V_{\omega} := \ker(H_{\omega} \rightarrow \mathbb{G}_m)$. There are morphisms of group schemes $G_{\omega} \rightarrow H_{\omega}$ and $U_{\omega} \rightarrow V_{\omega}$ compatible with the semi-direct product structure

$$\iota : G_{\omega} = \mathbb{G}_m \ltimes U_{\omega} \rightarrow H_{\omega} = \mathbb{G}_m \ltimes V_{\omega}. \quad (3)$$

Fix an embedding $\sigma : k \rightarrow \mathbb{C}$. Let $\Omega = \sum_{a \in \mu_N \cup \{0\}} \frac{dz}{z - \sigma(a)} e_a$. Paths $\mathbf{p}$ in $\mathbb{C}^* - \mu_N$ provide complex points $I(\mathbf{p}) = \sum_{n=0}^{\infty} \int_{\mathbf{p}} \Omega^n$ in Π with coordinates given by Chen's iterated integrals so that $I(\mathbf{p}$ followed by $\mathbf{q}) = I(\mathbf{q})I(\mathbf{p})$. At punctured points $x \in \{0, \infty\} \cup \mu_N$ one may use the tangential base points to extend the above construction to paths beginning or ending at x . This is a regularization procedure of divergent integrals with the counter terms prescribed by the tangent vectors. For example, one may define the straight line path from 1_0 (tangent vector 1 at 0) to $(-1)_1$ (see [3, p. 85]). This gives rise to $\text{dch}(\sigma)$ whose coefficient of $e_0^{s_1-1} e_{\zeta_1} \dots e_0^{s_n-1} e_{\zeta_n}$ is $(-1)^n Li_{s_1, \dots, s_n}(\sigma(1/\zeta_1), \sigma(\zeta_1/\zeta_2), \dots, \sigma(\zeta_{n-1}/\zeta_n))$ if $(s_1, \zeta_1) \neq (1, 1)$. Note that $\text{dch}(\sigma)$ is the unique group-like element in $\mathbb{Q}\langle\langle e_{\zeta} \rangle\rangle_{\zeta \in \mu_N \cup \{0\}}$ whose coefficients of e_0 and e_1 are both 0 with that property.

3 Standard relations in $\mathcal{MPV}(w, 4)$

Fix $N = 4$ throughout this section. In what follows we shall describe a process suggested by Deligne to verify that the standard relations cannot produce all possible $\mathbb{Q}$ -linear relations in $\mathcal{MPV}(w, 4)$.

Observe that the vector space freely generated by basis vectors corresponding to regularized MPVs of weight 3 and level 4 is dual to the degree 3 part of the free associative algebra $\mathbb{Q}\langle\langle e_0, e_\zeta \rangle\rangle_{\zeta \in \mu_4}$. One expects that $\mathcal{MPV}(w, 4)_{w \geq 1}$ is a weighted polynomial algebra with the following number of generators in low weights: 2, 1, 2, and 3 (in weight 1 to 4). The reason is that $\iota(\text{Lie } U_\omega)$ should be a Lie algebra freely generated by one element in each degree so that the dimension of degree n part is given by $\frac{1}{n} \sum_{d|n} \mu(n/d) 2^d - \delta_{1,n}$ (see [2, Ch. II, §3 Thm. 2]) which is the number of binary Lyndon words of length n when $n > 1$.

If one takes the space $\mathcal{MPV}(3, 4)$ modulo the products of MPVs in lower weights one should get a quotient space of dimension 2 and one knows its dimension ≤ 2 . However, one can only prove the bound 3 by the standard relations as follows. Consider the subspace generated by the following elements:

I. For each MPV of weight 1 and MPV of weight 2, the linear combination of weight 3 MPVs expressing their shuffle product by Chen's iterated integrals. This gives 5×25 elements.

II. Same for the stuffle product (quasi-shuffle as called by Hoffman) corresponding to the coproduct Δ_* in [7]. Plainly, these elements are linear combination of weight 3 MPVs expressing their stuffle products by series expansions (1). This gives 4×20 elements.

III. Distribution relations in weight 3: expressing the coefficient of a “convergent” monomial in e_0 and e_α ($\alpha = \pm 1$) as a multiple of the sum of the coefficients of the monomials deduced from it by replacing e_α by e_β with $\beta^2 = \alpha$. There are 12 “convergent” monomials to consider: $(e_0 \text{ or } e_{-1})(e_0 \text{ or } e_1 \text{ or } e_{-1})(e_1 \text{ or } e_{-1})$.

IV. Regularized distribution relations in weight 3 (see [7, Prop. 2.26], change $\sigma^{n/d} = 1$ to $\sigma^d = 1$). There are 6 of these.


All of these elements from I to IV can be put together to form a 223×125 matrix. By MAPLE one can verify that it only has rank 122. Moreover, it is not hard to find three linearly independent relations over $\mathbb{Q}$ which produce three vectors in the degree 2 part of the Lie algebra denoted by $\mathfrak{dmr}\mathfrak{d}_0$ in [7]. In weight 4 the same procedure produces eight vectors instead of three which means one needs five more relations besides the standard ones.

4 Octahedral symmetry of $\mathbb{P}^1 - (\{0, \infty\} \cup \mu_4)$

The Riemann sphere $\mathbb{P}^1$ punctured at $\{0, \infty, \pm 1, \pm i\}$ clearly possesses an octahedral symmetry. However, one should use a system of tangent vectors stable (up to multiplication by roots of unity) by the octahedral group to use this symmetry: one should use the straight line path from 1_0 to $(-2)_1$. Define the $\mathbb{C}$ -linear map $\rho : \mathbb{Q}\langle\langle e_0, e_\zeta \rangle\rangle_{\zeta \in \mu_4} \rightarrow \mathbb{Q}\langle\langle e_0, e_\zeta \rangle\rangle_{\zeta \in \mu_4}$ such that

$$\rho : e_0 \rightarrow e_1 \rightarrow e_i \rightarrow e_0, \quad e_\infty \rightarrow e_{-1} \rightarrow e_{-i} \rightarrow e_\infty.$$

Let $0 < \varepsilon < 1/3$ and C_ε be the path $A_1 \dots A_6$ in the complex plane shown in the right picture, where A_2, A_4 and A_6 are the quarter circles of radii ε , 2ε and 2ε , respectively, oriented clockwise. Then by the property of iterated integrals



$$1 = \int_{C_\varepsilon} \sum_{n=0}^{\infty} \Omega^{\circ n} = \int_{A_6} \sum_{n=0}^{\infty} \Omega^{\circ n} \int_{A_5} \sum_{n=0}^{\infty} \Omega^{\circ n} \dots \int_{A_1} \sum_{n=0}^{\infty} \Omega^{\circ n}. \quad (4)$$

Replacing the straight path $1_0 \rightarrow (-1)_1$ by the straight path $1_0 \rightarrow (-2)_1$ changes $\text{dch}(\sigma)$ to $\text{dch}'(\sigma) = \exp((\log 2)e_1) \text{dch}(\sigma)$. So regularized integral over the path A_1 followed by A_2 gives $I = \exp(-2\pi i e_1/4) \text{dch}'(\sigma)$. Thus (4) yields $\rho^2(I)\rho(I)I = 1$, i.e.,

$$\exp(-2Li_1(i)e_0)\rho^2(\text{dch}(\sigma))\exp(-2Li_1(i)e_i)\rho(\text{dch}(\sigma))\exp(-2Li_1(i)e_1)\text{dch}(\sigma) = 1. \quad (5)$$

By comparing the coefficient of $e_2e_1^2$ in (5) and using Fact (*) one finally arrives at (2).

We now consider the Lie algebra $\mathcal{G} := \iota(\text{Lie } U_\omega)$ which is the image of the map in (3), equipped with Ihara's Lie bracket $\{ , \}$ (see [4, (5.13.6)]). One can use the additional octahedral relation to find the correct number of generators of $\mathcal{G}$ in degrees up to 3:

$$\begin{aligned} \text{deg 1: } & v_1 = 2e_{-1} + 2e_1 + e_{-i} + e_i \\ \text{deg 2: } & v_2 = [e_0, e_i] - [e_0, e_{-i}] + [e_1, e_i] + [e_{-i}, e_1] + [e_{-i}, e_i] \\ \text{deg 3: } & \{v_1, v_2\} \text{ and } v_3, \end{aligned}$$

where v_3 is obtained by MAPLE and is too long to write down here. Define on the Lie algebra an involution $\sigma : e_0 \leftrightarrow e_1, e_i \leftrightarrow e_{-i}, e_{-1} \leftrightarrow e_\infty$ in the octahedral group. Then one can check by MAPLE that $v + \sigma(v) = 0$ for all the above generators.

Next we look at weight 4 case. We find that the upper bound produced by the standard relations is only $d(4, 4) \leq 21$ [8, Table 2]. By exactly the same approach as in weight 3 one can verify that all the five missing relations can be produced by applying the octahedral symmetry. For the five octahedral relations one can use the vanishing of the coefficients of $e_{-i}e_0^2e_{-i}$, $e_{-i}e_0^2e_{-1}$, $e_{-i}e_0^2e_i$, $e_{-i}e_0^2e_1$, and $(e_{-i}e_0)^2$ in (5). Moreover, a process similar to the one sketched in §3 yields exactly three generators in the space of the degree 4 part of the Lie algebra $\mathcal{G}$:

$$\text{deg 4: } \{v_1, v_3\}, \{v_1, \{v_1, v_2\}\}, \text{ and } v_4$$

where v_4 is a complicated new vector obtained by MAPLE. This verifies up to degree 4 the following fact: $\mathcal{G}$ is a Lie algebra freely generated by one generator in each degree.

Remark 1. For arbitrary N the set $\{0, \infty\} \cup \mu_N$ possesses the dihedral symmetry. However, when we apply this in $N = 4$ cases (weight 3 or 4) all the relations produced are already known to us by standard relations. This should be true in general (cf. [5, Thm. 4.1]).

5 Level p and p^2 cases ($p \geq 5$ a prime)

Let $p \geq 5$ be prime throughout this section. From Prop. 1 we see that $d(2, p) \leq (p+1)^2/4$. We will improve this bound in Theorem 1.

Recall that $\mathcal{G} = \iota(\text{Lie } U_\omega)$. By [4, 6.13] one may safely replace $\mathcal{G}_N^{(\ell)}$ by $\mathcal{G}$ throughout [5]. In particular, Goncharov's results can now be used to study the structure of Galois Lie algebra $\mathcal{G}_{\bullet, \bullet}(\mu_N) = \sum_{w \geq 1, l \geq 1} \mathcal{G}_{-w, -l}(\mu_N)$ graded by the weight w and depth l . The weight filtration is defined via the action of $\mathbb{G}_m$ and the depth filtration comes from the descending central series of $\ker f_*$, where $f_* : \pi_1(\mathbb{G}_m - \mu_N, x) \rightarrow \pi_1(\mathbb{G}_m)$ for any base point x is induced from the embedding $\mathbb{G}_m - \mu_N \hookrightarrow \mathbb{G}_m$.

Theorem 1. *Let $p \geq 5$ be a prime. Then*

$$d(2, p) \leq \frac{(5p+7)(p+1)}{24}. \quad (6)$$

If Grothendieck's period conjecture (see [4, 5.27(c)]) is true then the equality holds and all the $\mathbb{Q}$ -linear relations in $\mathcal{MPV}(2, p)$ follow from the standard relations.

Proof. Let $\{ , \}$ be the Lie bracket in $\mathcal{G}_{\bullet, \bullet}(\mu_N)$. Now in depth 1 one has $\dim \mathcal{G}_{-2, -1}(\mu_p) = (p-1)/2$ (see [5, Theorem 2.1]). By [5, Cor. 2.16]

$$\dim \mathcal{G}_{-2, -2}(\mu_p) = \frac{(p-1)(p-5)}{12}. \quad (7)$$

For any positive integer N define

$$\beta_N : \bigwedge^2 \mathcal{G}_{-2, -1}(\mu_N) \longrightarrow \mathcal{G}_{-2}(\mu_N), \quad a \wedge b \longmapsto \{a, b\}. \quad (8)$$

Then

$$\dim(\ker \beta_p) = \frac{1}{2} \frac{p-1}{2} \left(\frac{p-1}{2} - 1 \right) - \frac{(p-1)(p-5)}{12} = \frac{p^2-1}{24}. \quad (9)$$

Let SR be the affine subgroup of Π defined by only those polynomial equations satisfied by the coefficients of $\text{dch}(\sigma)$ which are deduced from the standard relations, plus “ $2\pi i = 0$ ”, as explained in [4, 5.22]. This is the group DMRD_0 studied by Racinet in [7, §3.2, Thm. I]. Its Lie algebra $\text{Lie } SR$ is graded by weight and depth, independent of the embedding by Prop. 4.1 of op. cit. and contains $\mathcal{G}$ by [4, 5.22]. Further, Goncharov shows [5, §7.7] that the standard relations provide a complete list of constraints on the diagonal part of the Lie algebra $\mathcal{G}$ in depth ≤ 2 , yielding $(\text{Lie } SR)_{m, m} = \mathcal{G}_{m, m}$ for $m = -1, -2$. Together with (9) this means in the proof of [4, 5.25] one can decrease the bound $D(2, p)$ by $(p^2-1)/24$ and arrive at the bound $(5p+7)(p+1)/24$.

Grothendieck's conjecture implies that $\mathcal{G} = \text{Lie } R$ where R is the affine subgroup of Π defined by all the polynomial equations satisfied by the coefficients of $\text{dch}(\sigma)$ plus “ $2\pi i = 0$ ”. Therefore one obtains both the equality in (6) and the statement about the completeness of the standard relations. This concludes the proof of the theorem. $\square$

Remark 2. (a) Notice that when $p \geq 11$ the vector space $\ker \beta_p$ contains a subspace isomorphic to the space of cusp forms of weight two on $X_1(p)$ which has dimension $(p-5)(p-7)/24$ (see [5, Lemma 2.3 & Theorem 7.8]). So it must contain another piece which has dimension $(p-3)/2$ by (9). What is this missing piece?

(b) When $w > 2$ it's not too hard to improve the bound of $d(w, p)$ given in [4, 5.24] by the same idea as used in the proof of the theorem (for example, decrease the bound by $(p^2-1)/24$). But they are often not the best.

Turn now to the case of level $N = p^2$. Notice that $\varphi(p^2)(\varphi(p^2)-2)/4$ is a trivial upper bound of $\dim \mathcal{G}_{-2, -2}(\mu_{p^2})$ (see [5, Theorem 2.1]). If one had any nontrivial upper bound of $\dim \mathcal{G}_{-2, -2}(\mu_{p^2})$ one would have a nontrivial lower bound of $\dim(\ker \beta_{p^2})$ where β_{p^2} is defined by (8). This would in turn provide an improved upper bound of $d(2, p^2)$ similar to Theorem 1. However, at present we can only show that

Theorem 2. *If $p \geq 5$ is a prime then $\ker \beta_{p^2} \neq 0$ and $d(2, p^2) < p^2(p-1)^2/4$.*

Proof. Fix a primitive p^2 th root of unity μ . Put $e(a) = e_{\mu^a}$ for all integer a . Define

$$g_{k, j} = e(pk + j) + e(p^2 - pk - j) + e(pj) + e(p^2 - pj)$$

for $0 \leq k < (p-1)/2$, $1 \leq j \leq p-1$, and for $k = (p-1)/2$, $1 \leq j \leq (p-1)/2$. One only needs to prove the following

Claim. Let $h = (p-3)/2$. Then one has

$$\begin{aligned} & \sum_{k=0}^h \sum_{l=k}^h \sum_{j=2}^{p-2} \{g_{k,1}, g_{l,j}\} + \sum_{k=0}^{h+1} \sum_{j=2}^{h+1} \{g_{k,1}, g_{h+1,j}\} + \sum_{k=0}^h \sum_{l=k+1}^h \sum_{j=2}^{p-2} \{g_{k,p-1}, g_{l,j}\} \\ & + \sum_{k=0}^h \sum_{j=2}^{h+1} \{g_{k,p-1}, g_{h+1,j}\} - \sum_{k=0}^h \sum_{l=k}^h \sum_{j=2}^{p-2} \{g_{k,j}, g_{l,p-1}\} - \sum_{k=0}^h \sum_{l=k}^h \sum_{j=2}^{p-2} \{g_{k,j}, g_{l+1,1}\} = 0. \end{aligned}$$

There are $h(2h+3)^2 = hp^2$ distinct terms on the left, each with coefficient ± 1 .

The proof of the claim is straight-forward by a little tedious change of indices and regrouping.

$$- \sum_{k=0}^h \sum_{l=k}^h \sum_{j=2}^{p-2} \{g_{k,j}, g_{l+1,1}\} = \sum_{k=0}^h \sum_{l=0}^k \sum_{j=2}^{p-2} \{g_{k+1,1}, g_{l,j}\} = \sum_{k=1}^{h+1} \sum_{l=0}^{k-1} \sum_{j=2}^{p-2} \{g_{k,1}, g_{l,j}\}.$$

Then the expression in the claim becomes

$$\begin{aligned} & \sum_{k=1}^h \sum_{l=0}^h \sum_{j=2}^{p-2} \{g_{k,1}, g_{l,j}\} + \sum_{l=0}^h \sum_{j=2}^{p-2} \{g_{0,1}, g_{l,j}\} + \sum_{l=0}^h \sum_{j=2}^{p-2} \{g_{h+1,1}, g_{l,j}\} + \sum_{k=0}^{h+1} \sum_{j=2}^{h+1} \{g_{k,1}, g_{h+1,j}\} \\ & + \sum_{k=0}^h \sum_{l=0}^h \sum_{j=2}^{p-2} \{g_{k,p-1}, g_{l,j}\} + \sum_{k=0}^h \sum_{j=2}^{h+1} \{g_{k,p-1}, g_{h+1,j}\} \\ & = \sum_{k=0}^{h+1} \sum_{l=0}^h \sum_{j=2}^{p-2} \{g_{k,1}, g_{l,j}\} + \sum_{k=0}^{h+1} \sum_{j=2}^{h+1} \{g_{k,1}, g_{h+1,j}\} \\ & + \sum_{k=0}^h \sum_{l=0}^h \sum_{j=2}^{p-2} \{g_{k,p-1}, g_{l,j}\} + \sum_{k=0}^h \sum_{j=2}^{h+1} \{g_{k,p-1}, g_{h+1,j}\} \end{aligned}$$

Let me use $\{a, b\} = \{e(a), e(b)\}$. By definition

$$\begin{aligned} \{g_{k,1}, g_{l,j}\} &= \{pk+1, pl+j\} + \{-pk-1, pl+j\} + \{p, pl+j\} + \{-p, pl+j\} \\ &+ \{pk+1, -pl-j\} + \{-pk-1, -pl-j\} + \{p, -pl-j\} + \{-p, -pl-j\} \\ &+ \{pk+1, pj\} + \{-pk-1, pj\} + \{p, pj\} + \{-p, pj\} \\ &+ \{pk+1, -pj\} + \{-pk-1, -pj\} + \{p, -pj\} + \{-p, -pj\} \\ &= \{pk+1, pl+j\} + \{p(p-k)-1, pl+j\} + \{p, pl+j\} + \{-p, pl+j\} \\ &+ \{pk+1, p(p-1-l)+p-j\} + \{p(p-k)-1, p(p-1-l)+p-j\} \\ &\quad + \{p, p(p-1-l)+p-j\} + \{-p, p(p-1-l)+p-j\} \\ &+ \{pk+1, pj\} + \{p(p-k)-1, pj\} + \{p, pj\} + \{-p, pj\} \\ &+ \{pk+1, p(p-j)\} + \{p(p-k)-1, p(p-j)\} + \{p, p(p-j)\} + \{-p, p(p-j)\} \end{aligned}$$

Then

$$\begin{aligned}
& \sum_{k=0}^{h+1} \sum_{l=0}^h \sum_{j=2}^{p-2} \{g_{k,1}, g_{l,j}\} = \sum_{k=0}^{h+1} \sum_{l=0}^h \sum_{j=2}^{p-2} \{pk+1, pl+j\} + \{p(p-k)-1, pl+j\} \\
& + \{pk+1, p(p-1-l)+j\} + \{p(p-k)-1, p(p-1-l)+j\} \\
& + \{p, pl+j\} + \{-p, pl+j\} + \{p, p(p-1-l)+j\} + \{-p, p(p-1-l)+j\} \\
& + 2\{pk+1, pj\} + 2\{p(p-k)-1, pj\} + 2\{p, pj\} + 2\{-p, pj\} \\
& = \sum_{k=0}^{h+1} \sum_{l=0, l \neq h+1}^{p-1} \sum_{j=2}^{p-2} \{pk+1, pl+j\} + \sum_{k=h+2}^p \sum_{l=0, l \neq h+1}^{p-1} \sum_{j=2}^{p-2} \{pk-1, pl+j\} \\
& + \sum_{k=0}^{h+1} \sum_{l=0, l \neq h+1}^{p-1} \sum_{j=2}^{p-2} (\{p, pl+j\} + \{-p, pl+j\}) + 2(h+1) \sum_{k=0}^{h+1} \sum_{j=2}^{p-2} \{pk+1, pj\} \\
& + 2(h+1) \sum_{k=h+2}^p \sum_{j=2}^{p-2} \{pk-1, pj\} + 2(h+2)(h+1) \sum_{j=2}^{p-2} (\{p, pj\} + \{-p, pj\})
\end{aligned}$$

Thus

$$\begin{aligned}
& \sum_{k=0}^{h+1} \sum_{l=0}^h \sum_{j=2}^{p-2} \{g_{k,1}, g_{l,j}\} + \sum_{k=0}^{h+1} \sum_{j=2}^{h+1} \{g_{k,1}, g_{h+1,j}\} \\
& = \sum_{k=0}^{h+1} \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} \{pk+1, pl+j\} + \sum_{k=h+2}^p \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} \{pk-1, pl+j\} \\
& + \frac{p+1}{2} \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} (\{p, pl+j\} + \{-p, pl+j\}) + p \sum_{k=0}^{h+1} \sum_{j=2}^{p-2} \{pk+1, pj\} \\
& + p \sum_{k=h+2}^p \sum_{j=2}^{p-2} \{pk-1, pj\} + \frac{p^2-1}{2} \sum_{j=2}^{p-2} (\{p, pj\} + \{-p, pj\})
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \sum_{k=0}^h \sum_{l=0}^h \sum_{j=2}^{p-2} \{g_{k,p-1}, g_{l,j}\} + \sum_{k=0}^h \sum_{j=2}^{h+1} \{g_{k,p-1}, g_{h+1,j}\} \\
& = \sum_{k=1}^{h+1} \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} \{pk-1, pl+j\} + \sum_{k=h+2}^{p-1} \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} \{pk+1, pl+j\} \\
& + \frac{p-1}{2} \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} (\{p, pl+j\} + \{-p, pl+j\}) + p \sum_{k=1}^{h+1} \sum_{j=2}^{p-2} \{pk-1, pj\} \\
& + p \sum_{k=h+2}^{p-1} \sum_{j=2}^{p-2} \{pk+1, pj\} + \frac{(p-1)^2}{2} \sum_{j=2}^{p-2} (\{p, pj\} + \{-p, pj\})
\end{aligned}$$

Altogether the expression in the claim is reduced to

$$\begin{aligned}
& \sum_{k=0}^{p-1} \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} \{pk+1, pl+j\} + \sum_{k=1}^p \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} \{pk-1, pl+j\} \\
& + p \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} \left(\{p, pl+j\} + \{-p, pl+j\} \right) + p \sum_{k=0}^{p-1} \sum_{j=2}^{p-2} \{pk+1, pj\} \\
& + p \sum_{k=1}^p \sum_{j=2}^{p-2} \{pk-1, pj\} + p(p-1) \sum_{j=2}^{p-2} \left(\{p, pj\} + \{-p, pj\} \right)
\end{aligned}$$

This is equal to

$$\begin{aligned}
& \sum_{k=0}^{p-1} \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} \left([pk+1, pl+j] - [p(k+l)+j+1, pl+j] + [p(k+l)+j+1, pk+1] \right) \\
& + \sum_{k=1}^p \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} \left([pk-1, pl+j] - [p(k+l)+j-1, pl+j] + [p(k+l)+j-1, pk+1] \right) \\
& + p \sum_{l=0}^{p-1} \sum_{j=2}^{p-2} \left([p, pl+j] - [p(l+1)+j, pl+j] + [p(l+1)+j, p] \right. \\
& \quad \left. + [-p, pl+j] - [p(l-1)+j, pl+j] + [p(l-1)+j, -p] \right) \\
& + p \sum_{k=0}^{p-1} \sum_{j=2}^{p-2} \left([pk+1, pj] - [p(j+k)+1, pj] + [p(j+k)+1, pk+1] \right) \\
& + p \sum_{k=1}^p \sum_{j=2}^{p-2} \left([pk-1, pj] - [p(j+k)-1, pj] + [p(j+k)-1, pk-1] \right) \\
& + p(p-1) \sum_{j=2}^{p-2} \left([p, pj] - [p(j+1), pj] + [p(j+1), p] \right. \\
& \quad \left. + [-p, pj] - [p(j-1), pj] + [p(j-1), -p] \right) = 0
\end{aligned}$$

after many cancellations. □

We have verified the claim in the above proof for $p = 5$ and $p = 7$ numerically. In fact, we find that $\dim(\ker \beta_{25}) = 5$ and $\dim(\ker \beta_{49}) = 35$. This implies that $d(2, 25) \leq 116$ and $d(2, 49) \leq 449$. Further, if Grothendieck's transcendence conjecture is true then the equalities hold. With MAPLE we have verified $d(2, 25) \leq 116$ and $d(2, 49) \leq 449$ by using the standard relations so that these relations imply all the others.

It is convenient to say a level N is *standard* if either (i) $N = 1, 2$ or 3 , or (ii) N is a prime power p^n ($p \geq 5$). Otherwise N is called *non-standard*. We prove in [8] the following

Theorem 3. *Suppose N is non-standard. Then the non-standard relations exist in $\mathcal{MPV}(3, N)$ for $4 \leq N \leq 12$ and in $\mathcal{MPV}(2, N)$ for $10 \leq N \leq 48$.*

Preliminary results suggest that the following

Conjecture 1. *If N is a standard level then the standard relations always provide the sharp bounds of $d(w, N)$, namely, all linear relations can be derived from the standard ones. If N is a non-standard level then the bound in [4, Cor. 5.25] is sharp and the non-standard relations exist in $\mathcal{MPV}(w, N)$ for all $w \geq 3$ (and in $\mathcal{MPV}(2, N)$ if $N \geq 10$).*

Acknowledgements. I would like to express my deepest gratitude to Prof. Deligne for patiently explaining [4] to me and suggesting to use octahedral symmetry in level four. This note was written when I was visiting the Institute for Advanced Study at Princeton whose financial support and hospitality is gratefully acknowledged.

References

- [1] J. Borwein, P. Lisonek, and P. Irvine, *An interface for evaluation of Euler sums*, available online at <http://oldweb.cecm.sfu.ca/cgi-bin/EZFace/zetaform.cgi>
- [2] N. Bourbaki, *Lie Groups and Lie Algebras Chapters 1-3*, Springer, 2004.
- [3] P. Deligne, *Le groupe fondamental de la droite projective moins trois points*, Galois groups over $\mathbb{Q}$, (Berkeley, CA, 1987), Springer, New York, 1989, p. 79–297.
- [4] P. Deligne and A. Goncharov, *Groupes fondamentaux motiviques de Tate mixte*, Ann. Scient. Éc. Norm. Sup., **38** (1)(2005), 1–56.
- [5] A. Goncharov, *The dihedral Lie algebras and Galois symmetries of $\pi_1^{(l)}(\mathbb{P}_1 - (\{0, \infty\} \cup \mu_N))$* , Duke Math. J. **110** (3)(2001), 397–487.
- [6] J. Vollinga, S. Weinzierl, *Numerical evaluation of multiple polylogarithms*, Computer Phys. Comm., **167** (3)(2005), 177–194.
- [7] G. Racinet, *Doubles mélanges des polylogarithmes multiples aux racines de l’unité*, Publ. Math. Inst. HautesEtudes Sci. **95** (2002), 185–231.
- [8] J. Zhao, *Standard relations of multiple polylogarithm values of at roots of unity*, preprint: arXiv:0707.1459v7.